

SUBORDINATION PROBLEMS OF ROBERTSON FUNCTIONS

LI-MEI WANG

ABSTRACT. In the present paper, we are concerned with subordination problems related to λ -Robertson function. The radii of λ -spirallikeness and starlikeness of λ -Robertson function are also determined.

1. INTRODUCTION AND MAIN RESULTS

Let $\mathbb{D}_r = \{z \in \mathbb{C} : |z| < r\}$ for $0 < r \leq 1$ and $\mathbb{D} = \mathbb{D}_1$ be the unit disc. Let \mathcal{A} be the family of functions f analytic in \mathbb{D} , and \mathcal{A}_1 be the subset of \mathcal{A} consisting of functions f which are normalized by $f(0) = f'(0) - 1 = 0$. A function $f \in \mathcal{A}$ is said to be subordinate to a function $F \in \mathcal{A}$ in \mathbb{D} (in symbols $f \prec F$ or $f(z) \prec F(z)$) if there exists an analytic function $\omega(z)$ on \mathbb{D} with $|\omega(z)| < 1$ and $\omega(0) = 0$, such that

$$f(z) = F(\omega(z))$$

in \mathbb{D} . When F is a univalent function, the condition $f \prec F$ is equivalent to $f(\mathbb{D}) \subseteq F(\mathbb{D})$ and $f(0) = F(0)$. Let

$$\mathcal{P}_\lambda = \{p \in \mathcal{A} : p(0) = 1, \operatorname{Re} e^{-i\lambda} p(z) > 0\}.$$

Here and hereafter we always suppose $-\pi/2 < \lambda < \pi/2$. Note that \mathcal{P}_λ is a convex and compact subset of \mathcal{A} which is equipped with the topology of uniform convergence on compact subsets of \mathbb{D} . Since \mathcal{P}_0 is the well-known Carathéodory class, we call \mathcal{P}_λ the *tilted Carathéodory class by angle* λ . Some characterizations and estimates of elements in \mathcal{P}_λ are known (for a short survey, see [11]).

For a function $f \in \mathcal{A}$, let

$$Q_f(z) = \frac{zf'(z)}{f(z)} \tag{1}$$

and

$$P_f(z) = 1 + \frac{zf''(z)}{f'(z)}.$$

It is worthwhile to note that

$$Q_f(z) + \frac{zQ'_f(z)}{Q_f(z)} = P_f(z).$$

Primary 30C45, Secondary 30C62, 30C75

Key words and Phrases: Robertson functions, spirallike functions, subordination theory, radii of spirallikeness and starlikeness.

These quantities are important for investigation of geometric properties of analytic functions. Next we will define two subclasses of analytic functions related to these two quantities.

A function $f \in \mathcal{A}_1$ is said to be a λ -*spirallike function* (denoted by $f \in \mathcal{SP}_\lambda$) if

$$Q_f \in \mathcal{P}_\lambda.$$

Note that \mathcal{SP}_0 is precisely the set of starlike functions normally denoted by \mathcal{S}^* . Spirallike functions were introduced and proved to be univalent by Špaček [10] in 1932. For general references about spirallike functions, see e.g. [2] or [1].

A function $f \in \mathcal{A}_1$ is said to be a λ -*Robertson function* if $zf'(z) \in \mathcal{SP}_\lambda$, i.e.

$$P_f \in \mathcal{P}_\lambda.$$

Let \mathcal{R}_λ denote the set of these functions. Note that \mathcal{R}_0 is precisely the set of convex functions sometimes denoted by \mathcal{K} . Convex functions have been the subject of numerous investigations, among which the following result was proved by MacGregor [6] in 1975.

Theorem A. *Let $f \in \mathcal{K}$, then the subordination relation*

$$\frac{zf'(z)}{f(z)} \prec \frac{zf_0'(z)}{f_0(z)}$$

holds, where $f_0(z) = z/(1-z)$.

We are interested in more general subordination problems related to λ -Robertson functions. For this purpose, we first introduce some specific functions for convenience.

A distinguished member of \mathcal{R}_λ is

$$f_\lambda(z) = \frac{(1-z)^{1-2e^{i\lambda}\cos\lambda} - 1}{2e^{i\lambda}\cos\lambda - 1}.$$

A simple calculation yields

$$Q_\lambda(z) = \frac{zf'_\lambda(z)}{f_\lambda(z)} = \frac{e^{2i\lambda}z}{1-z-(1-z)^{1+e^{2i\lambda}}}, \quad (2)$$

$$P_\lambda(z) = 1 + \frac{zf''_\lambda(z)}{f'_\lambda(z)} = \frac{1+e^{2i\lambda}z}{1-z}$$

and

$$Q_\lambda(z) + \frac{zQ'_\lambda(z)}{Q_\lambda(z)} = P_\lambda(z). \quad (3)$$

In [4], Kim and Srivastava posed the open problem which is an extension of Theorem A whether

$$\frac{zf'(z)}{f(z)} \prec \frac{zf'_\lambda(z)}{f_\lambda(z)}$$

holds for $f \in \mathcal{R}_\lambda$ with general λ . In other words, if

$$q(z) + \frac{zq'(z)}{q(z)} \prec P_\lambda(z) \quad (4)$$

in \mathbb{D} , then whether

$$q \prec Q_\lambda$$

holds in \mathbb{D} for general λ ? Relation (4) is a kind of Briot-Bouquet differential subordinations which have a surprising number of important applications in the theory of univalent functions. Many sources and references are given in [7].

In the present paper, we solve the above problem in a restricted disc and obtain the radii of spirallikeness and starlikeness for Robertson functions as well.

Theorem 1. *Let $q \in \mathcal{A}$ with $q(0) = 1$ satisfy the differential subordination*

$$q(z) + \frac{zq'(z)}{q(z)} \prec P_\lambda(z) \quad (5)$$

and $R_1(\lambda)$ be defined by

$$R_1(\lambda) = \sup\{r < 1 : Q_\lambda(rz) \prec P_\lambda(rz) \text{ in } \mathbb{D}\}. \quad (6)$$

Then

$$q(z) \prec Q_\lambda(z)$$

in $|z| < R_1(\lambda)$.

By the discussion in Section 1, we can deduce the following corollary immediately from Theorem 1.

Corollary 1. *Let $f \in \mathcal{R}_\lambda$, then*

$$\frac{zf'(z)}{f(z)} \prec \frac{zf'_\lambda(z)}{f_\lambda(z)}$$

in $|z| < R_1(\lambda)$, where $R_1(\lambda)$ is given in (6).

Remark 1. The radius of λ -spirallikeness of λ -Robertson functions is at least $R_1(\lambda)$, since

$$\frac{zf'(z)}{f(z)} \prec \frac{zf'_\lambda(z)}{f_\lambda(z)} \prec P_\lambda(z)$$

in $\mathbb{D}_{R_1(\lambda)}$ for any $f \in \mathcal{R}_\lambda$.

Remark 2. For $0 \leq r < 1$, let

$$\begin{aligned} \psi_\lambda(r) &= \max_{|z|=1} |P_\lambda^{-1}(Q_\lambda(rz))/r| \\ &= \max_{|z|=1} \left| \frac{1}{r} \frac{mrz - 1 + (1 - rz)^m}{1 - (1 - rz)^m} \right| \end{aligned}$$

where $m = 1 + e^{2i\lambda}$. $\psi_\lambda(r)$ is an increasing function defined on $[0, 1]$ with $\psi_\lambda(0) = 0$. By the definition of subordination, $R_1(\lambda)$ defined in (6) could be expressed in terms of ψ_λ :

$$R_1(\lambda) = \sup\{r < 1 : \psi_\lambda(r) < 1\} = \psi_\lambda^{-1}(1).$$

Theorem 2. *Let $q \in \mathcal{A}$ with $q(0) = 1$ satisfy the differential subordination*

$$q(z) + \frac{zq'(z)}{q(z)} \prec P_\lambda(z), \quad (7)$$

then

$$q(R_2 z) \prec P_0(z)$$

in \mathbb{D} , where

$$R_2 := R_2(\lambda) = \frac{2}{\sqrt{4 + 2\sqrt{3}|\sin(2\lambda)|}}. \quad (8)$$

Corollary 2. *The radius of starlikeness of λ -Robertson functions is at least $R_2(\lambda)$ given in (8).*

Note that $R_1(0) = R_2(0) = 1$, thus both Corollary 1 and Corollary 2 imply Theorem A. Note also that in [1], Ahuja and Silverman posed the problem to find the radius of starlikeness for all λ -Robertson functions. Corollary 2 implies that this radius is at least

$$R_2 = \min\{R_2(\lambda) : -\pi/2 < \lambda < \pi/2\} = R_2(\pi/4) = \sqrt{3} - 1 \approx 0.732.$$

Libera and Ziegler in [5] have shown that the radius of close-to-convexity for all λ -Robertson functions is approximately 0.99097524 and the radius of convexity is $\sqrt{2}/2$.

2. PROOFS OF RESULTS

In order to obtain our main results, the following lemmas are required.

Lemma 1 ([11]). *Let $p \in \mathcal{P}_\lambda$, then we have*

$$|p(z) - A(r)| \leq B(r),$$

where

$$A(r) = \frac{1 + r^2 e^{2i\lambda}}{1 - r^2}, \quad B(r) = \frac{2r \cos \lambda}{1 - r^2}$$

and $r = |z| < 1$. Equality holds if and only if $p(z) = P_\lambda(xz)$ with $|x| = 1$.

Lemma 2 ([7], Lemma 2.2d). *Let $g(z)$ and $h(z)$ be in \mathcal{A} with $g(0) = h(0)$. If $g \not\prec h$ in \mathbb{D} , then there exist two points z_0 with $|z_0| < 1$ and η_0 with $|\eta_0| = 1$ and $s \geq 1$ such that*

$$g(\mathbb{D}_{|z_0|}) \subset h(\mathbb{D}),$$

$$g(z_0) = h(\eta_0)$$

and

$$z_0 g'(z_0) = s \eta_0 h'(\eta_0).$$

The next lemma is due to Nunokawa [8]. We only quote the relevant part.

Lemma 3 ([8]). *Let $p(z) \in \mathcal{A}$ satisfy $p(0) = 1$ and $p(z) \neq 0$ in \mathbb{D} . If there exists a point $z_0 \in \mathbb{D}$ such that $\operatorname{Re} p(z) > 0$ in $|z| < |z_0|$ and $p(z_0) = ia$ where $a \in \mathbb{R} \setminus \{0\}$, then*

$$\frac{z_0 p'(z_0)}{p(z_0)} = ik$$

where $k \geq (a + 1/a)/2$ if $a > 0$ and $k \leq -(a + 1/a)/2$ if $a < 0$.

Proof of Theorem 1. For simplicity, we let $R = R_1(\lambda)$ and $p(z) = q(z) + zq'(z)/q(z)$, thus $p \in \mathcal{P}_\lambda$. If $q(z) \not\prec Q_\lambda(z)$ in $|z| < R$, then Lemma 2 implies the existences of z_0 with $|z_0| < R$, η_0 with $|\eta_0| = R$ and $s \geq 1$ such that

$$\begin{aligned} q(\mathbb{D}_{|z_0|}) &\subset Q_\lambda(\mathbb{D}), \\ q(z_0) &= Q_\lambda(\eta_0), \\ z_0 q'(z_0) &= s \eta_0 Q'_\lambda(\eta_0). \end{aligned} \tag{9}$$

Thus in view of (3), (5) and (9), we have

$$\begin{aligned} p(z_0) &= q(z_0) + \frac{z_0 q'(z_0)}{q(z_0)} \\ &= Q_\lambda(\eta_0) + s \frac{\eta_0 Q'_\lambda(\eta_0)}{Q_\lambda(\eta_0)} \\ &= s P_\lambda(\eta_0) + (1 - s) Q_\lambda(\eta_0). \end{aligned} \tag{10}$$

Therefore (10) and Lemma 1 show that

$$\begin{aligned} |p(z_0) - A(R)| &= |s(P_\lambda(\eta_0) - A(R)) - (1 - s)(Q_\lambda(\eta_0) - A(R))| \\ &\geq |s(P_\lambda(\eta_0) - A(R))| - (s - 1)|Q_\lambda(\eta_0) - A(R)| \\ &\geq sB(R) - (s - 1)B(R) \\ &= B(R) \end{aligned}$$

which contradicts with $p(z) \in \mathcal{P}_\lambda$. Therefore we get the assertion. \square

Proof of Theorem 2. For simplicity, we let $R = R_2(\lambda)$ and $p(z) = q(z) + zq'(z)/q(z)$, thus $p \in \mathcal{P}_\lambda$. If $q(Rz) \not\prec P_0(z)$ in \mathbb{D} , it follows from $q(0) = 1$ that there exists a point $z_0 \in \mathbb{D}$ such that $\operatorname{Re} q(Rz) > 0$ for $|z| < |z_0|$ and $q(Rz_0) = ia$ where $a \in \mathbb{R} \setminus \{0\}$, then by Lemma 3, we have

$$\frac{Rz_0 q'(Rz_0)}{q(Rz_0)} = ik,$$

where $k \geq (a + 1/a)/2$ if $a > 0$ and $k \leq -(a + 1/a)/2$ if $a < 0$. Therefore

$$p(Rz_0) = q(Rz_0) + \frac{Rz_0 q'(Rz_0)}{q(Rz_0)} = ia + ik,$$

which implies $p(Rz_0) \in \Omega$ since $|a + k| \geq \sqrt{3}$, where $\Omega = \{it, |t| \geq \sqrt{3}\}$. Next we will show that

$$p(\mathbb{D}_R) \cap \Omega = \emptyset,$$

which contradicts the above assertion. Since $p \in \mathcal{P}_\lambda$, it is sufficient to prove for functions $P_\lambda(z)$. Suppose that there is a point $z_1 \in \mathbb{D}$ such that $P_\lambda(z_1) = it_0$ with $|t_0| \geq \sqrt{3}$, then a simple calculation gives that

$$z_1 = \frac{it_0 - 1}{it_0 + e^{2i\lambda}}.$$

Hence

$$|z_1|^2 = \frac{t_0^2 + 1}{t_0^2 + 1 - 2t_0 \sin(2\lambda)} \geq \frac{4}{4 + 2\sqrt{3}|\sin(2\lambda)|} = R$$

since $|t_0| \geq \sqrt{3}$. Therefore $P_\lambda(\mathbb{D}_R) \cap \Omega = \emptyset$. The proof is completed. \square

Acknowledgements

The author is grateful to Professor Toshiyuki Sugawa for his guidance and useful discussions during the preparation of this paper.

REFERENCES

- [1] O. P. Ahuja and H. Silverman, *A survey on spiral-like and related function classes*, Math. Chronicle **20** (1991), 39–66.
- [2] P. L. Duren, *Univalent functions*, Springer-Verlag, New York, 1983.
- [3] A. W. Goodman, *Univalent Functions*, 2 vols., Mariner Publishing Co. Inc., 1983.
- [4] Y. C. Kim and H. M. Srivastava, *Some subordination properties for spirallike functions*, Appl. Math. Comput. **203** (2008), 838–842.
- [5] R. J. Libera and M. R. Ziegler, *Regular functions $f(z)$ for which $zf'(z)$ is α -spiral*, Trans. Amer. Math. Soc. **166** (1972), 361–370.
- [6] T. H. MacGregor, *A subordination for convex functions of order a* , J. London Math Soc. (2), **9** (1975), 530–536.
- [7] S. Miller and P. Mocanu, *Differential Subordinations: Theory and Applications*, Series on Monographs and Textbooks in Pure and Applied Mathematics (No. 225), Marcel Dekker, New York and Basel, 2000.
- [8] M. Nunokawa, *On properties of non-Carathéodory functions*, Proc. Japan Acad. Ser. A Math. Sci. **68**, (1992), no. 6, 152–153.
- [9] M. S. Robertson, *Univalent functions $f(z)$ for which $zf'(z)$ is spirallike*, Michigan Math. J. **16** (1969), 97–101.
- [10] L. Špaček, *Contribution à la théorie des fonctions univalentes*, Cas. Mat. Fys **62** (1932), 12–19.
- [11] L.-M. Wang, *The tilted Carathéodory class and its applications*, preprint (to appear in J. Korean Math. Soc.).

Division of Mathematics

Graduate School of Information Sciences

Tohoku University, Sendai

980-8579 JAPAN

e-mail: rime@ims.is.tohoku.ac.jp

wangmabel@163.com